

**WELCOME  
TO**



**Drx Notes**

**biochemistry | Chapter-5**


# Nucleic acids

---

- Definition, purine and pyrimidine bases
- Components of nucleosides and nucleotides with examples
- Structure of DNA (Watson and Crick model), RNA and their functions


## Purine and Pyrimidine

- Purines and pyrimidine are both organic compounds that take part in the synthesis of DNA and RNA, therefore they are called as the building blocks of the genetic material – DNA and RNA.
- They are nitrogenous bases that make up the two different nucleotides in DNA and RNA.



- Purine is a heterocyclic aromatic organic compound composed of a pyrimidine ring fused with imidazole ring. It comprises adenine and guanine as nucleases.
- It consists of two hydrogen-carbon rings and four nitrogen atoms.
- The melting point of purine is 214 °C. Catabolism results in the production of uric acid

**Example-Purines (adenine and guanine) are two-carbon nitrogen ring base.**



- Pyrimidine is a heterocyclic aromatic organic compound that is composed of carbon and hydrogen.
- It comprises cytosine, thymine, uracil as nucleobases.
- It consists of one hydrogen-carbon ring and two nitrogen atoms.
- The melting point of pyrimidine is 20-22 °C. Catabolism produces carbon dioxide, beta-amino acids and ammonia

**Example - pyrimidines (cytosine and thymine) are one-carbon nitrogen ring bases.**

## Nucleoside and Nucleotide


1. Nitrogenous Base: They comprise pyrimidine or purine base. DNA contains adenine (A), guanine (G), cytosine (C) and thymine (T) whereas RNA contains adenine, guanine, cytosine and uracil (U).
2. Sugar: A nucleotide comprises a pentose sugar. DNA (Deoxyribonucleic acid) contains deoxyribose sugar and RNA (Ribonucleic acid) contains a ribose sugar.
  - A Nitrogenous base attached with the sugar is called "Nucleoside".
3. Phosphate: Phosphate is associated with the sugar of nucleoside by an ester bond with the 5thC hydroxyl group. Nucleotides at least contain one phosphate group.
  - A nucleoside is a purine or a pyrimidine nucleobase with a pentose sugar component, which is either ribose or deoxyribose.

Therefore:

**nucleoside = nucleobase + ribose or deoxyribose**

- A nucleoside is a glycoside formed from the hydrolysis of nucleic acid.
- In a nucleoside, the anomeric carbon is attached to the N9 of a purine (or to the N1 of a pyrimidine) by a glycosidic bond.


### Nucleoside structure:



- Nucleotides are organic molecules consisting of a nucleoside and a phosphate.
- They serve as monomeric units of the nucleic acid **polymers – deoxyribonucleic acid** and **ribonucleic acid**, both of which are essential biomolecules within all life-forms on Earth.
- A molecule consisting of a nitrogen-containing base (adenine, guanine, thymine, or cytosine in DNA; adenine, guanine, uracil, or cytosine in RNA), a phosphate group, and a sugar (deoxyribose in DNA; ribose in RNA).


***nucleotide = nucleoside + phosphate group***

### Nucleotide structure:



### Watson and Crick model of DNA

- DNA as an acidic substance present in the nucleus was first identified by Frederich Meischer in 1869. He named it as 'nucleon'. Due to technical limitations in isolating such a long polymer intact the elucidation of structure of DNA remained elusive for a long period of time.
- It was only in 1953 that James Watson and Francis Crick proposed the very simple but famous double helix model for the structure of DNA.
- The main opposition was base pairing between the two strands of polynucleotide chains.



### ***Structure of DNA double helix***

**The salient features of double helix structure of DNA are as follows:**

- It is made up of two polynucleotide chains.
- The two chains have antiparallel polarity if one has polarities and the second chain must have polarity.
- The base into strands is paired through hydrogen bond forming base pairs. Adenine forms two hydrogen bonds with thymine from opposite strands and vice versa.
- Similarly guanine forms three H bonds with cytosine. As a result, purine comes opposite to pyrimidine.
- Because of this approximate a uniform distance between the two strengths of The Helix occurs.
- The two chains are called in a right-handed fashion. Pitch of the helix is and there are roughly 10bp in each turn.
- The plane of one base pair is stacked over the other in a double helix. This confirms stability of the helical structure.

**Note: The proposition of a double helix structure for DNA and its simplicity in explaining the genetic implication become revolutionary.**

### **Function of DNA and RNA:**

#### **RNA:**

- You can think of an RNA molecule, as a disposable copy of a segment of DNA, a working copy of a single gene.
- RNA has many functions, but most RNA molecules are involved in protein synthesis only.
- RNA controls the assembly of amino acids into proteins.

- Each type of RNA molecule specializes in a different aspect of this job. The three main types of RNA are messenger RNA, ribosomal RNA, and transfer RNA.

**DNA has two functions:**

1. **Hold information on how to make proteins**
2. **Make more DNA**

**The main function of DNA is to store genetic information.**

- The information that DNA stores is how, when, and where to make protein.
- The second function of DNA is simply to make more DNA; this is called replication.



# Drx Notes